

AD-A275 109



AGE

Form Approved  
GSA No. 27-1130

(2)

1. AGENCY USE

December 27, 1993

3. REPORT TYPE AND DATES COVERED

Reprint

4. TITLE AND SUBTITLE

Large-Scale Coronal Temperature and Density  
Distributions, 1984-1992

5. FUNDING NUMBERS

PE 61102F

PR 2311

TA G3

WU 27

6. AUTHOR(S)

M. Guhathakurta\*, R.R. Fisher\*\*, R.C. Altrock

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Phillips Lab/GPSS  
29 Randolph Road  
Hanscom AFB, MA 01731-30108. PERFORMING ORGANIZATION  
REPORT NUMBER

PL-TR-93-2254

9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES)

10. SPONSORING MONITORING  
AGENCY REPORT NUMBERJAN 6 1994  
S B L

\*Laboratory for Atmospheric and Space Physics, University of  
Colorado, Boulder, CO 80303 \*\*NASA/Goddard Space Flight Center, Code 682, Greenbelt  
MD 20771

Reprinted from The Astrophysical Journal, 414: L145-L148, 1993 September 10

Approved for public release; Distribution unlimited

In this Letter we characterize the temperature and the density structure of the corona utilizing spectrophotometric observations at different heights but at the same latitude during the descending phase of cycle 21 through the ascending phase of cycle 22. The data include ground-based intensity observations of the green (Fe xiv  $\lambda 5303$ ) and red (Fe x  $\lambda 6374$ ) coronal forbidden lines, photospheric magnetographs from the National Solar Observatory, Kitt Peak, and synoptic maps of white-light K-coronal polarized brightness.  $pB$ , from the High Altitude Observatory. A determination of plasma temperature  $T$  can be estimated from the intensity ratio Fe x/Fe xiv (where  $T$  is inversely proportional to the ratio), since both emission lines come from ionized states of Fe, and the ratio is only weakly dependent on density. Distributions of the electron temperature from the line ratio and the polarized brightness which yields electron density of the corona during the descending and the ascending phases of solar cycles 21 and 22 are presented. These data refer to structures of the corona which are relatively large scale, having a temporal coherence of at least two or more synoptic rotation periods, such as the streamer belts, the individual helmet streamers, and the larger coronal holes. We observe that there is a large-scale organization of the inferred coronal temperature distribution that is associated with the large-scale structures in the solar magnetic fields; this organization tends to persist through most of the magnetic activity cycle. This distribution differs in spatial and temporal characterization from the traditional picture of sunspot and active region evolution over the range of sunspot cycles, which are manifestations of the small-scale, strong magnetic field regions. For example, during 1987-1990 the active region latitudes drift equatorward, whereas bright coronal features in white light and the inferred temperature structures drift poleward.

14. SUBJECT TERMS

Solar corona, Solar temperature, Solar density

15. NUMBER OF PAGES

5

16. PRICE CODE

17. SECURITY CLASSIFICATION  
OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION  
OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION  
OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

SAR

**Best  
Available  
Copy**

## LARGE-SCALE CORONAL TEMPERATURE AND DENSITY DISTRIBUTIONS, 1984-1992

M. GUHATHAKURTA<sup>1</sup>

Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303

R. R. FISHER

NASA Goddard Space Flight Center, Code 682, Greenbelt, MD 20771

AND

R. C. ALTROCK

Phillips Laboratory (AFSC), Geophysics Directorate, National Solar Observatory, Sacramento Peak, Sunspot, NM 88349

Received 1992 November 5, accepted 1993 June 24

## ABSTRACT

In this *Letter* we characterize the temperature and the density structure of the corona utilizing spectrophotometric observations at different heights but at the same latitude during the descending phase of cycle 21 through the ascending phase of cycle 22. The data include ground-based intensity observations of the green (Fe XIV  $\lambda 5303$ ) and red (Fe X  $\lambda 6374$ ) coronal forbidden lines, photospheric magnetographs from the National Solar Observatory, Kitt Peak and synoptic maps of white-light K-coronal polarized brightness,  $pB$ , from the High Altitude Observatory. A determination of plasma temperature  $T$  can be estimated from the intensity ratio Fe X/Fe XIV (where  $T$  is inversely proportional to the ratio), since both emission lines come from ionized states of Fe, and the ratio is only weakly dependent on density. Distributions of the electron temperature from the line ratio and the polarized brightness which yields electron density of the corona during the descending and the ascending phases of solar cycles 21 and 22 are presented. These data refer to structures of the corona which are relatively large scale, having a temporal coherence of at least two or more synoptic rotation periods, such as the streamer belts, the individual helmet streamers, and the larger coronal holes. We observe that there is a large-scale organization of the inferred coronal temperature distribution that is associated with the large-scale structures in the solar magnetic fields; this organization tends to persist through most of the magnetic activity cycle. This distribution differs in spatial and temporal characterization from the traditional picture of sunspot and active region evolution over the range of sunspot cycles, which are manifestations of the small-scale, strong magnetic field regions. For example, during 1987-1990 the active region latitudes drift equatorward, whereas bright coronal features in white light and the inferred temperature structures drift poleward.

*Subject headings:* Sun: corona

## 1. INTRODUCTION

Knowledge of the basic conditions of temperature and density in the corona is fundamental to our understanding of the dominant physical processes that drive the corona and its solar-terrestrial connections via the interplanetary medium. However, very few accurate measurements have been made of the temperature in the inner corona. Recently, Guhathakurta et al. (1992) and Guhathakurta & Altrock (1992) determined the temperature structure of the inner corona utilizing cotemporal observations of the X-ray and ultraviolet intensities, the coronal green line Fe XIV, and the coronal red line Fe X during the total solar eclipse of 1988 March, and estimated average temperatures during the period 1984-1992, respectively. These studies provide a systematic method for the study of the altitudinal variation of the temperature and density as a function of the phase of the solar cycle.

The present study considers only the largest scale structure of the solar corona. The photometric data were sampled at a daily rate, and in the creation of the synoptic charts of distributions a sample size of  $6^\circ$  of solar latitude was used. This means that this study was band-limited to features with a spatial scale not less than  $12^\circ$  of latitude in width as seen at the limb, and at

least 2 days' rotation ( $\sim 26^\circ$ ) in longitude. Scale sizes set by these limits are approximately  $0.2$  and  $0.4 R_\odot$ , respectively, and include helmet streamer, streamer belt, and larger scale coronal hole structures.

Daily observations of the solar corona are made at the National Solar Observatory at Sacramento Peak with the Photoelectric Coronal Photometer (Smartt 1982). These observations have been made in the Fe lines at 6374 and 5303 Å which are formed at approximate temperatures of 1 and 2 MK, respectively (Altrock 1990). The 1:1 entrance aperture is scanned daily around the limb at  $1.15 R_\odot$ .

Since 1980, the imaging K-coronameter Mark III (Fisher et al. 1984) of the High Altitude Observatory at Mauna Loa Solar Observatory in Hawaii has measured the distribution of coronal polarized brightness in white light as a function of height and azimuth around the limb of the Sun almost daily.

A set of synoptic observations of the longitudinal component of the photospheric magnetic field has been collected by the National Solar Observatory at Kitt Peak, and this set is used to provide a sense of the spatial and temporal evolution of the magnetic field of the Sun during the period of this study.

## 2. INTERPRETATION OF OBSERVED LINE PROFILES

In order to interpret the observed intensities in the Fe lines, we derive theoretical intensity profiles of these lines to be com-

<sup>1</sup> Summer faculty at NASA/GFSC, Greenbelt, MD

94-00273



94 1 5 010



persisted for several rotations, it would be seen in this display as a series of repetitive bright areas extending along the x-axis. This repetitive pattern would have a temporal frequency of about 6.7 per year due to the frequency of sampling the data. An example of rotational modulation can be seen in the white-light and green line records especially near the declining phase (1984) and minimum (1986) of the cycle. We have used 16 levels of color to represent white-light data ranging from  $0.1$  to  $22 \times 10^{-8} I_{\odot}$  (disk center brightness). In this first strip of plot (top), black represents any data point below  $1.38 \times 10^{-8} I_{\odot}$ , the range of colors from dark to light green is  $(1.38-4.13) \times 10^{-8} I_{\odot}$ , beige is  $(4.13-4.5) \times 10^{-8} I_{\odot}$ , light blue to dark blue is  $(4.5-9.63) \times 10^{-8} I_{\odot}$ , purple to pink is  $(9.63-13.75) \times 10^{-8} I_{\odot}$ , yellow to red is  $(13.75-19.25) \times 10^{-8} I_{\odot}$ , and gray to white is  $(19.25-22) \times 10^{-8} I_{\odot}$ . The total intensity and latitude extent of the white-light corona is modulated at the frequency of the solar magnetic cycle. In the white-light plot, it is seen that as the cycle declines in 1984 toward the minimum at 1986, the white-light  $pB$  drops in amplitude (green is low  $pB$ ) to a minimum and then toward the middle of 1987 begins to increase toward maximum values of  $pB$  which are displayed in pink, yellow, red near the maximum of this current cycle, 1989-1990.

The reader is reminded that there are two timescales visualized as the eye scans the x-axis. These two periodic variations result from the modulation by solar rotation and the variation associated with the solar magnetic activity cycle. Rotational modulation of the observed signal occurs at a frequency of about 6.7 cycles  $\text{yr}^{-1}$  and magnetic activity cycle modulation occurs at the rate of  $\sim 11$  cycle  $\text{yr}^{-1}$ . Solar cycle modulation of the observed quantities is easily seen in the Fe x (red line) and Fe xiv (green line) emission-line records which are plotted in the next to bottom and bottom panels of Figure 2, respectively. Since these lines are controlled largely by collisional excitation, we expect them to reflect the variation associated with approximately the square of the local electron density. The active regions are easy to pick out by inspection, since an increase in the local electron density of a factor of 3 will account for approximately an order of magnitude increase in line brightness. The solar magnetic cycle modulation of the emission-line corona by the active regions are clearly seen in these two data sets with the onset of the new cycle regions starting in mid-1988. In the red line plot, black represents any data point below  $0.75 \times 10^{-6} B_{\odot}$ , the range of colors from dark to light green is  $(0.75-2.25) \times 10^{-6} B_{\odot}$ , beige is  $(2.25-3.0) \times 10^{-6} B_{\odot}$ , light blue to dark blue is  $(3.0-5.25) \times 10^{-6} B_{\odot}$ , purple to pink is  $(5.25-7.5) \times 10^{-6} B_{\odot}$ , yellow to red is  $(7.5-10.5) \times 10^{-6} B_{\odot}$ , and gray to white is  $(10.5-12) \times 10^{-6} B_{\odot}$ . Similarly, in the green line plot, black represents any data point below  $3.13 \times 10^{-6} B_{\odot}$ , dark to light green is  $(3.13-9.38) \times 10^{-6} B_{\odot}$ , beige is  $(9.38-12.5) \times 10^{-6} B_{\odot}$ , light blue to dark blue is  $(12.5-21.9) \times 10^{-6} B_{\odot}$ , purple to pink is  $(21.9-31.13) \times 10^{-6} B_{\odot}$ , yellow to red is  $(31.13-43.75) \times 10^{-6} B_{\odot}$ , and gray to white is  $(43.75-50) \times 10^{-6} B_{\odot}$ .

The photospheric magnetic field as detected by the NSO longitudinal magnetograph is plotted on the strip of data which is second from the top. In this case strong fields of opposite polarity are outlined in white, gray and black, green, while the weak fields are shaded red and blue. The 1984-1985 data have been rescaled to be consistent with the data available from 1985 to 1992, and both the rotational and the magnetic cycle modulation of the signal as a function of time are evident.

The inferred coronal temperature from the Fe x Fe xiv ratio is shown in the middle strip of data in Figure 2. Relatively cool regions in the corona are colored green ( $\sim 1.25-1.35$  MK), the intermediate regions ( $\sim 1.45-1.6$  MK) are colored beige, blue and the pink, yellow, and red regions are areas where the inferred temperature is higher (near  $1.8-2.1$  MK). The most prominent features of the data display are the two bands of higher temperature (pink, yellow, and red) material which are seen at relatively high latitudes on either side of the solar equator. The average temperature in these bands is found to be  $1.85 \pm 0.10$  MK, which is about 500,000 K hotter than the average temperature ( $1.32 \pm 0.07$  MK) at the poles during solar minimum and 300,000 K hotter than the mean equatorial temperature ( $1.57 \pm 0.11$  MK) determined for the period of this study.

At the time near solar minimum, at 1986, there is relatively little variation of latitude as a function of time in the high-temperature regions. Near the maximum of the magnetic activity cycle there is a drift of the poleward boundary toward higher latitudes, first seen in the southern hemisphere, and then, about half a year later, a similar enlargement of the high-temperature region of the corona is seen to migrate toward the north polar region.

Inspection of the latitudes of the high-temperature bands yields evidence that is consistent with the location of large-scale weak-field structure in the solar magnetic field. For example, the rapid poleward spread in the high-temperature band during the 1987-1989 period of rising solar activity occurred while small-scale manifestations of activity, such as sunspots and active regions, were slowly drifting toward the equator. Further inspection of the combined photospheric magnetic field, and of the temperature, indicates that regions of higher temperature tend to be located over positions where weak photospheric fields of opposite sign are in the process of being transported toward the pole of rotation (Guhathakurta & Fisher 1992). Although not yet demonstrated by a quantitative statistical analysis, we find that prominences and bright features in the quiet corona (both related to large-scale magnetic fields) migrated toward high latitudes during 1987-1989 (Hundhausen 1993) in a manner consistent with the inferred high-temperature bands.

From the compilation of H $\alpha$  synoptic charts McIntosh (1979) uses the dark disk filaments (prominences seen against the bright solar disk) in mapping the complete patterns of large-scale solar magnetic polarity. These long, persistent filaments which mark most of the patterns reveal lines of polarity reversal also known as "neutral" lines (NLs). Prominences, regions of cool, dense material at coronal heights in the vicinity of these magnetic neutral lines, are thought to be supported against solar gravity by a surrounding magnetic field structure with a spatial scale comparable to the prominence length ( $\sim 0.3 R_{\odot}$ ; Allen 1973). The NLs that underlie the polar crown of filaments are the longest on the solar surface, interconnecting the two hemispheres. The so-called rush to the poles by the polar crown filaments is part of a process of poleward migration by a series of large-scale patterns that merge into a single feature in the year or two before maximum. This merger creates the nearly continuous chain of filaments known as the polar crown (McIntosh 1979). We observe a striking similarity in the latitudinal distributions of these polar crown filaments (McIntosh 1922; Cliver et al. 1993) during 1984-1992, and the inferred high-temperature coronal regions.

The polarized brightness distribution in Figure 2, to first order, reflects the total electron content of the corona independent of temperature. This electron content can be seen as the sum of two components, each with a distinct association with detected photospheric fields, thermal characteristics, and latitude variation with time over the cycle.

The first component is the fraction of the corona associated with the large-scale magnetic field and the boundary between weak opposite fields that is best seen near solar minimum. The white-light corona, in this case, is composed of large-scale bright persistent structures with relatively high latitudes and moderate to high density (Fig. 2). The bright features typically correspond to the centers of coronal helmet streamers (Guhathakurta et al. 1992), which are thought to have closed magnetic field lines that extend outward up to a height of about a solar radius. Enhanced inferred temperatures are seen at these latitudes.

The second component of the total variation in density of the corona over the period of study is that associated with the small-scale magnetic field and with high field strength regions of spot groups and active regions. The latitudinal variation with time is the same as that for the spot groups, drifting toward the solar equator as time advances from the maximum of the cycle. These dense coronal structures do not demonstrate elevated temperatures and to the first order seem to have a characteristic temperature which is independent of density (Guhathakurta & Fisher 1992).

In this analysis, we have used several data sets, for the purposes of comparisons, that were measured at different heights in the solar atmosphere. No effort was made to extrapolate these data to the same height. For example, the green and red line data were measured at  $1.15 R_{\odot}$ , the white-light data at  $1.3 R_{\odot}$ , and the photospheric magnetic field data at  $1 R_{\odot}$ . However, we find the spatial correspondence between green line bright spots and the active regions in the magnetic field data to be in close agreement, indicating that we are still observing the same features with these data sets. The latitudes of features seen on the solar disk, such as sunspots, active regions, etc., and the apparent latitudes of coronal features seen in limb observations made in white light have been studied by Hundhausen (1993), and he draws the conclusion that it is unlikely that the high-latitude coronal bright features ( $50^{\circ}$ – $60^{\circ}$ ) are the projection of structures at active region latitudes ( $25^{\circ}$ – $45^{\circ}$ ). Thus the inferred high-latitude, high-temperature band obtained by using emission lines, which are approximately proportional to  $n^2$ , is not likely to be the result of projection effects of the rotation of structures at active region latitudes.

## 5. CONCLUSIONS

The following conclusions were drawn

1. There is an organization of the large-scale coronal temperature distribution which is associated with the large-scale structure in the solar magnetic field. Generally this organization takes the form of two zonal bands each about  $20^{\circ}$  of solar latitude in width, where the latitude boundaries are located around  $50^{\circ}$  N and  $50^{\circ}$  S.

2. These structures tend to persist through most of the solar activity cycle. Near the maximum of the sunspot cycle there was a poleward expansion of these zones while small-scale manifestations of activity such as sunspots and active regions were slowly drifting toward the equator.

3. The temperature of these zones at the reference height of  $1.15 R_{\odot}$  is about 300,000 K hotter than the mean temperature inferred for the corona near the equator of the Sun and about 500,000 K hotter than the mean temperature inferred at the polar regions during solar minimum.

4. Recent analyses show that the high-temperature material of these zones tends to lie over regions where magnetograph observations indicate a change of polarity of weak large-scale magnetic fields (Guhathakurta & Fisher 1992).

5. Finally, the latitude distributions of the small-scale aspects of the solar activity as observed in sunspots and active regions, and reflected in the green and red line data and the longitudinal component of the magnetic field in Figure 2 (the so-called butterfly diagram), are significantly different from the distributions of the large-scale features as observed in white-light data and the inferred temperature structures. The latitudinal distributions of the large- and small-scale structures change in very different ways during the ascent to maximum phase (1987–1990) of the solar cycle. During this phase the active region latitudes drift equatorward, whereas bright coronal features in white light and the inferred temperature structures drift poleward.

This research was supported by a NASA summer faculty fellowship to one of us (M. G.), who would like to thank all the individuals in the SDAC group at GSFC for all the assistance with facilities and analysis that they provided. Observations at NSO/SP were obtained by Evans Facility observers under the supervision of Lou B. Gilliam, Chief Observer. K-coronameter data were supplied to us by D. Sime of the HAO. Some data reduction support was provided by Timothy W. Henry (NSO/SP) and Vic Tisone (NCAR/HAO). We would like to thank J. W. Harvey (NSO/KP) for providing the synoptic magnetogram data. Finally, we would like to thank the referee for many useful suggestions.

## REFERENCES

- Allen, C. W. 1973, *Astrophysical Quantities* (3d ed., London: Athlone).
- Altrock, R. C. 1990, in *Climate Impact of Solar Variability* (NASA CP-3086), 287.
- 1993, private communication.
- Arnaud, M., & Rothenflug, R. 1985, *A&AS*, 60, 425.
- Cliver, E. W., St. Cyr, O. C., Howard, R. A., & McIntosh, P. S. 1993, in *Proc. 23d Cosmic Ray Conference* (Calgary), in press.
- Fisher, R. R., McCabe, M., Mickey, L., Seagraves, P., & Sime, D. G. 1984, *ApJ*, 280, 873.
- Guhathakurta, M., & Altrock, R. C. 1992, in *ASP Conf. Ser. Vol. 27, The Solar Cycle* (San Francisco: ASP), 395.
- Guhathakurta, M., & Fisher, R. R. 1992, *BAAS*, 24, 4, 1254.
- Guhathakurta, M., Rottman, G. J., Fisher, R. R., Orrall, F. Q., & Altrock, R. 1992, *ApJ*, 388, 633.
- Hundhausen, A. J. 1993, *J. Geophys. Res.*, in press.
- Mason, H. E. 1975, *MNRAS*, 170, 651.
- McIntosh, P. S. 1979, UAG Report 70, World Data Center A for Solar Terrestrial Physics, Boulder.
- 1992, in *ASP Conf. Ser. Vol. 27, The Solar Cycle* (San Francisco: ASP), 14.
- Smartt, R. N. 1982, *Proc. SPIE*, 231, 442.




FIG. 2. Evolution of solar corona as seen at (a) green line intensity (bottom strip) in units of  $10^{-12} B_{\odot}$  (total brightness of the solar disk; Allen 1973), (b) red line intensity in units of  $10^{-12} B_{\odot}$ , (c) the temperature estimate in MK, from the line ratio (middle strip) of the inner corona at  $1.15 R_{\odot}$ , (d) the photospheric magnetic field from Kitt Peak, and (e) the coronal white light (top strip) at  $1.5 R_{\odot}$  from Mauna Loa, Hawaii, in units of  $10^{-12} I_{\odot}$  (disk center brightness of the solar disk; Allen 1973), for Carrington rotations 1743-1860 for heliographic latitudes  $-90^{\circ}$  to  $90^{\circ}$ . The color scale is linear and is plotted against each panel with their respective maxima and minima.